

Families of Rationally Connected Varieties

February 1, 2008

Tom Graber

Mathematics Department, Harvard University,
1 Oxford st., Cambridge MA 02138, USA
graber@math.harvard.edu

Joe Harris

Mathematics Department, Harvard University,
1 Oxford st., Cambridge MA 02138, USA
harris@math.harvard.edu

Jason Starr

Mathematics Department, M.I.T.,
Cambridge MA 02139, USA
jstarr@math.mit.edu

Contents

1	Introduction	2
1.1	Statement of results	2
2	Preliminary definitions and constructions	5
2.1	The first construction	6
2.2	The second construction	9
3	Proof of the main theorem	10
3.1	The proof in case $B = \mathbb{P}^1$	10
3.2	The proof for arbitrary curves B	13
4	An example	14

1 Introduction

1.1 Statement of results

We will work throughout over the complex numbers, so that the results here apply over any algebraically closed field of characteristic 0.

Recall that a smooth projective variety X is said to be *rationally connected* if two general points $p, q \in X$ can be joined by a chain of rational curves. In case $\dim(X) \geq 3$ this turns out to be equivalent to the a priori stronger condition that for any finite subset $\Gamma \subset X$ there is a smooth rational curve $C \subset X$ containing Γ and having ample normal bundle.

Rationally connected varieties form an important class of varieties. In dimensions 1 and 2 rational connectivity coincides with rationality, but the two notions diverge in higher dimensions and in virtually every respect the class of rationally connected varieties is better behaved. For example, the condition of rational connectivity is both open and closed in smooth proper families; there are geometric criteria for rational connectivity (e.g. any smooth projective variety with negative canonical bundle is rationally connected, so we know in particular that a smooth hypersurface $X \subset \mathbb{P}^n$ of degree d will be rationally connected if and only if $d \leq n$), and there are, at least conjecturally, numerical criteria for rational connectivity (see Conjecture 1.6 below). In this paper we will prove a conjecture of Kollár, Miyaoka and Mori that represents one more basic property of rational connectivity (also one not shared by rationality): that if $X \rightarrow Y$ is a morphism with rationally connected image and fibers, then the domain X is rationally connected as well. This will be a corollary of our main theorem:

Theorem 1.1 *Let $f : X \rightarrow B$ be a morphism from a smooth projective variety to a smooth projective curve over $\mathbb{C}$. If the general fiber of f is rationally connected, then f has a section.*

Since this is really a statement about the birational equivalence class of the morphism f , we can restate it in the equivalent form

Theorem 1.2 *If K is the function field of a curve over $\mathbb{C}$, any rationally connected variety X defined over K has a K -rational point.*

In this form, the theorem directly generalizes Tsen's theorem, which is exactly this statement for X a smooth hypersurface of degree $d \leq n$ in

projective space $\mathbb{P}^n$ (or more generally a smooth complete intersection in projective space with negative canonical bundle). It would be interesting to know if in fact rationally connected varieties over other C_1 fields necessarily have rational points.

As we indicated, one basic corollary of our main theorem is

Corollary 1.3 *Let $f : X \rightarrow Y$ be any dominant morphism of varieties. If Y and the general fiber of f are rationally connected, then X is rationally connected.*

Proof. We can assume (in characteristic 0, at least) that X and Y are smooth projective varieties. Let p and q be general points of X . We can find a smooth rational curve $C \subset Y$ joining $f(p)$ and $f(q)$; let $X' = f^{-1}(C)$ be the inverse image of C in X . By Theorem 1.1, there is a section D of X' over C . We can then connect p to q by a chain of rational curves in X' in three stages: connect p to the point $D \cap X_p$ of intersection of D with the fiber X_p of f through p by a rational curve; connect $D \cap X_p$ to $D \cap X_q$ by D , and connect $D \cap X_q$ to q by a rational curve in X_q . □

There is a further corollary of Theorem 1.1 based on a construction of Campana and Kollár–Miyaoka–Mori: the *maximal rationally connected fibration* associated to a variety X (see [Ca], [K] or [KMM]). Briefly, the maximal rationally connected fibration associates to a variety X a (birational isomorphism class of) variety Z and a rational map $\phi : X \rightarrow Z$ with the properties that

- the fibers X_z of ϕ are rationally connected; and conversely
- almost all the rational curves in X lie in fibers of ϕ : for a very general point $z \in Z$ any rational curve in X meeting X_z lies in X_z .

The variety Z and morphism ϕ are unique up to birational isomorphism, and are called the *mrc quotient* and *mrc fibration* of X , respectively. They measure the failure of X to be rationally connected: if X is rationally connected, Z is a point, while if X is not uniruled we have $Z = X$. As observed in Kollár ([K], IV.5.6.3), we have the following Corollary:

Corollary 1.4 *Let X be any variety and $\phi : X \rightarrow Z$ its maximal rationally connected fibration. Then Z is not uniruled.*

Proof. Suppose that Z were uniruled, so that through a general point $z \in Z$ we could find a rational curve $C \subset Z$ through z . By Corollary 1.3, the inverse image $\phi^{-1}(C)$ will be rationally connected, which means that every point of the fiber X_z will lie on a rational curve not contained in X_z , contradicting the second defining property of mrc fibrations. □

There are conjectured numerical criteria for a variety X to be either uniruled or rationally connected. They are

Conjecture 1.5 *Let X be a smooth projective variety. Then X is uniruled if and only if $H^0(X, K_X^m) = 0$ for all $m > 0$.*

and

Conjecture 1.6 *Let X be a smooth projective variety. Then X is rationally connected if and only if $H^0(X, (\Omega_X^1)^{\otimes m}) = 0$ for all $m > 0$.*

For each of these conjectures, the “only if” part is known, and straightforward to prove; the “if” part represents a very difficult open problem (see for example [K], IV.1.12 and IV.3.8.1). As another consequence of our main theorem, we have an implication:

Corollary 1.7 *Conjecture 1.5 implies Conjecture 1.6*

Proof. Let X be any smooth projective variety that is not rationally connected; assuming the statement of Conjecture 1.5, we want to show that $H^0(X, (\Omega_X^1)^{\otimes m}) \neq 0$ for some $m > 0$. Let $\phi : X \rightarrow Z$ be the mrc fibration of X . By hypothesis Z has dimension $n > 0$, and by Corollary 1.4 Z is not uniruled. If we assume Conjecture 1.5, then, we must have a non-zero section $\sigma \in H^0(Z, K_Z^m)$ for some $m > 0$. But the line bundle K_Z^m is a summand of the tensor power $(\Omega_Z^1)^{\otimes nm}$, so we can view σ as a global section of that sheaf; pulling it back via ϕ , we get a nonzero global section of $(\Omega_X^1)^{\otimes nm}$. □

Acknowledgments. We would like to thank Johan deJong, János Kollár and Barry Mazur for many conversations, which were of tremendous help to us.

2 Preliminary definitions and constructions

We will be dealing with morphisms $\pi : X \rightarrow B$ satisfying a number of hypotheses, which we collect here for future reference. In particular, for the bulk of this paper we will deal with the case $B \cong \mathbb{P}^1$; we will show in section 3.2 below both that the statement for $B \cong \mathbb{P}^1$ implies the full Theorem 1.1 and, as well, how to modify the argument that follows to apply to general B .

Hypotheses 2.1 $\pi : X \rightarrow B$ is a nonconstant morphism of smooth connected projective varieties over $\mathbb{C}$, with $B \cong \mathbb{P}^1$. For general $b \in B$, the fiber $X_b = \pi^{-1}(b)$ is rationally connected of dimension at least 2.

Now suppose we have a class $\beta \in N_1(X)$ having intersection number d with a fiber of the map π . We have then a natural morphism

$$\varphi : \overline{M}_{g,0}(X, \beta) \rightarrow \overline{M}_{g,0}(B, d)$$

defined by composing a map $f : C \rightarrow X$ with π and collapsing components of C as necessary to make the composition πf stable.

Definition 2.2 Let $\pi : X \rightarrow B$ be a morphism satisfying 2.1, and let $f : C \rightarrow X$ be a stable map from a nodal curve C of genus g to X with class $f_*[C] = \beta$. We say that f is *flexible* relative to π if the map $\varphi : \overline{M}_{g,0}(X, \beta) \rightarrow \overline{M}_{g,0}(B, d)$ is dominant at the point $[f] \in \overline{M}_{g,0}(X, \beta)$; that is, if any neighborhood of $[f]$ in $\overline{M}_{g,0}(X, \beta)$ dominates a neighborhood of $[\pi f]$ in $\overline{M}_{g,0}(B, d)$.

Now, it's a classical fact that the variety $\overline{M}_{g,0}(B, d)$ has a unique irreducible component whose general member corresponds to a flat map $f : C \rightarrow B$ (see for example [C] and [H] for a proof). Since the map $\varphi : \overline{M}_{g,0}(X, \beta) \rightarrow \overline{M}_{g,0}(B, d)$ is proper, it follows that if $\pi : X \rightarrow B$ admits a flexible curve then φ will be surjective. Moreover, $\overline{M}_{g,0}(B, d)$ contains points $[f]$ corresponding to maps $f : C \rightarrow B$ with the property that every irreducible component of C on which f is nonconstant maps isomorphically via f to B . (For example, we could simply start with d disjoint copies $C_1, \dots, C_d$ of B (with f mapping each isomorphically to B) and identify $d + g - 1$ pairs of points on the C_i , each pair lying over the same point of B).

Proposition 2.3 *If $\pi : X \rightarrow B$ is a morphism satisfying 2.1 and $f : C \rightarrow X$ a flexible stable map, then π has a section.*

Our goal in what follows, accordingly, will be to construct a flexible curve $f : C \rightarrow X$ for an arbitrary $\pi : X \rightarrow B$ satisfying 2.1.

2.1 The first construction

To manufacture our flexible curve, we apply two basic constructions, which we describe here. (These constructions, especially the first, are pretty standard: see for example section II.7 of [K].) We start with a basic lemma:

Lemma 2.4 *Let C be a smooth curve and E any vector bundle on C ; let n be any positive integer. Let $p_1, \dots, p_N \in C$ be general points and $\xi_i \subset E_{p_i}$ a general one-dimensional subspace of the fiber of E at p_i ; let E' be the sheaf of rational sections of E having at most a simple pole at p_i in the direction ξ_i and regular elsewhere. For N sufficiently large we will have*

$$H^1(C, E'(-q_1 - \dots - q_n)) = 0$$

for any n points $q_1, \dots, q_n \in C$.

Proof. To start with, we will prove simply that $H^1(C, E') = 0$. Since this is an open condition, it will suffice to exhibit a particular choice of points p_i and subspaces ξ_i that works. Denoting the rank of E by r , we take $N = mr$ divisible by r and choose m points $t_1, \dots, t_m \in C$. We then specialize to the case

$$\begin{aligned} p_1 = \dots = p_r = t_1; & \quad \xi_1, \dots, \xi_r \text{ spanning } E_{t_1} \\ p_{r+1} = \dots = p_{2r} = t_2; & \quad \xi_{r+1}, \dots, \xi_{2r} \text{ spanning } E_{t_2} \end{aligned}$$

and so on. In this case we have $E' = E(t_1 + \dots + t_m)$, which we know has vanishing higher cohomology for sufficiently large m .

Given this, the statement of the lemma follows: to begin with, choose any $g + n$ points $r_1, \dots, r_{g+n} \in C$. Applying the argument thus far to the bundle $E(-r_1 - \dots - r_{g+n})$, we find that for N sufficiently large we will have $H^1(C, E'(-r_1 - \dots - r_{g+n})) = 0$. But now for any points $q_1, \dots, q_n \in C$ we have

$$q_1 + \dots + q_n = r_1 + \dots + r_{g+n} - D$$

for some effective divisor D on C . It follows then that

$$\begin{aligned} h^1(C, E'(-q_1 - \dots - q_n)) &= h^1(C, E'(-r_1 - \dots - r_{g+n})(D)) \\ &\leq h^1(C, E'(-r_1 - \dots - r_{g+n})) \\ &= 0 \end{aligned}$$

□

The relevance of this to our present circumstances will perhaps be made clear by the following:

Lemma 2.5 *Let X be a smooth projective variety, C and $C' \subset X$ two nodal curves meeting at points $p_1, \dots, p_\delta$; suppose C and C' are smooth with distinct tangent lines at each point p_i . Let $D = C \cup C'$ be the union of C and C' ; and let $N_{C/X}$ and $N_{D/X}$ be the normal sheaves of C and D in X . We have then an inclusion of sheaves*

$$0 \rightarrow N_{C/X} \rightarrow N_{D/X}|_C$$

identifying the sheaf of sections of $N_{D/X}|_C$ with the sheaf of rational sections of $N_{C/X}$ having at most a simple pole at p_i in the normal direction determined by $T_{p_i}C'$. Moreover, if $\tilde{D} \subset \text{Spec } \mathbb{C}[\epsilon]/(\epsilon^2) \times X$ is a first-order deformation of D in X corresponding to a global section $\sigma \in H^0(N_{D/X})$, then $\tilde{D}$ smooths the node of D at p_i if and only if the restriction $\sigma|_U$ of σ to a neighborhood U of p in C is not in the image of $N_{C/X}$.

Now suppose $\pi : X \rightarrow B$ is a morphism satisfying our basic hypotheses 2.1, and $C \subset X$ a smooth, irreducible curve of genus g . For a general point $p \in C$, let $X_p = \pi^{-1}(\pi(p))$ be the fiber of π through p . By hypothesis, X_p is a smooth, rationally connected variety, so that we can find a smooth rational curve $C' \subset X_p$ meeting C at p (and nowhere else) with arbitrarily specified tangent line at p , and having ample normal bundle $N_{C'/X}$.

Choose a large number of general points $p_1, \dots, p_\delta \in C$, and for each i let $C_i \subset X_{p_i}$ be such a smooth rational curve, with $T_{p_i}C_i$ a general tangent line to X_{p_i} at p_i . Combining the preceding two lemmas, we see that for δ sufficiently large, the normal bundle $N_{C'/X}$ of the union $C' = C \cup (\cup C_i)$ will be generated by its global sections; in particular, by Lemma 2.5 there will be a smooth deformation $\tilde{C}$ of C' . Moreover, for any given n we can choose the number δ large enough to ensure that $H^1(C, N_{C'/X}|_C(-r_1 - \dots - r_{g+n})) = 0$ for some $g+n$ points $r_1, \dots, r_{g+n} \in C$; it follows that $H^1(\tilde{C}, N_{\tilde{C}/X}(-r_1 - \dots - r_{g+n})) = 0$ for some $r_1, \dots, r_{g+n} \in \tilde{C}$ and hence that

$$H^1(\tilde{C}, N_{\tilde{C}/X}(-q_1 - \dots - q_n)) = 0$$

for any n points on $\tilde{C}$.

The process of taking a curve $C \subset X$, attaching rational curves in fibers and smoothing to get a new curve $\tilde{C}$, is our first construction. It has the properties that

1. the genus g of the new curve $\tilde{C}$ is the same as the genus of the curve C we started with;
2. the degree d of $\tilde{C}$ over B is the same as the degree of C over B ;
3. the branch divisor of the composite map $\tilde{C} \hookrightarrow X \rightarrow B$ is a small deformation of the branch divisor of $C \hookrightarrow X \rightarrow B$; and again,
4. for any n points $q_1, \dots, q_n \in \tilde{C}$ we have $H^1(\tilde{C}, N_{\tilde{C}/X}(-q_1 - \dots - q_n)) = 0$

Here is one application of this construction. Suppose we have a smooth curve $C \subset X$ such that the projection $\mu : \pi|_C : C \rightarrow B$ is simply branched—that is, the branch divisor of μ consists of $2d + 2g - 2$ distinct points in B —and such that each ramification point $p \in C$ of μ is a smooth point of the fiber X_p . Applying our first construction with $n = 2d + 2g - 2$, we arrive at another smooth curve $\tilde{C}$ that is again simply branched over B , with all ramification occurring at smooth points of fibers of π . But now the condition that $H^1(\tilde{C}, N_{\tilde{C}/X}(-q_1 - \dots - q_n)) = 0$ applied to the $n = 2d + 2g - 2$ ramification points of the map $\tilde{\mu} : \tilde{C} \rightarrow B$ says that if we pick a normal vector v_i to $\tilde{C}$ at each ramification point p_i of $\tilde{\mu}$ we can find a global section of the normal bundle $N_{\tilde{C}/X}$ with value v_i at p_i . Moreover, since ramification occurs at smooth points of fibers of π , for any tangent vectors w_i to B at the image points $\pi(p_i)$ we can find tangent vectors $v_i \in T_{p_i}X$ with $d\pi(v_i) = w_i$. It follows that *as we deform the curve $\tilde{C}$ in X , the branch points of $\tilde{\mu}$ move independently*. A general deformation of $\tilde{C} \subset X$ thus yields a general deformation of $\tilde{\mu}$ —in other words, the curve $\tilde{C}$ is flexible. We thus make the

Definition 2.6 Let $\pi : X \rightarrow B$ be as in 2.1, and let $C \subset X$ be a smooth curve such that the projection $\mu : \pi|_C : C \rightarrow B$ is simply branched. If each ramification point $p \in C$ of μ is a smooth point of the fiber X_p containing it, we will say the curve C is *pre-flexible*.

In these terms, we have established the

Lemma 2.7 *Let $\pi : X \rightarrow B$ be as in 2.1. If X admits a pre-flexible curve, the map π has a section.*

Remark. Note that we can extend the notion of pre-flexible and the statement of Lemma 2.7 to stable maps $f : C \rightarrow X$: we say that such a map is preflexible is the composition πf is simply branched, and for each ramification point p of πf the image $f(p)$ is a smooth point of the map π , the statement of Lemma 2.7 holds.

2.2 The second construction

Our second construction is a very minor modification of the first. Given a family $\pi : X \rightarrow B$ as in 2.1 and a smooth curve $C \subset X$, we pick a general fiber X_b of π and two points $p, q \in C \cap X_b$. We then pick a rational curve $C_0 \subset X_b$ with ample normal bundle in X_b , passing through p and q and not meeting C elsewhere. We also pick a large number N of other general points $p_i \in C$ and rational curves $C_i \subset X_{p_i}$ in the corresponding fibers, meeting C just at p_i and having general tangent line at p_i . Finally, we let $C' = C \cup C_0 \cup (\cup C_i)$ be the union, and $\tilde{C}$ a smooth deformation of C' (as before, if we pick N large enough, the normal bundle $N_{C'/X}$ will be generated by global sections, so smoothings will exist). This process, starting with the curve $C \subset X$ and arriving at the new curve $\tilde{C}$, is our second construction. It has the properties that

1. the degree d of $\tilde{C}$ over B is the same as the degree of C over B ;
2. the genus of the new curve $\tilde{C}$ is one greater than the genus of the curve C we started with;
3. for any n points $q_1, \dots, q_n \in \tilde{C}$ we have $H^1(\tilde{C}, N_{\tilde{C}/X}(-q_1 - \dots - q_n)) = 0$; and
4. the branch divisor of the composite map $\tilde{C} \hookrightarrow X \rightarrow B$ has two new points: it consists of a small deformation of the branch divisor of $C \hookrightarrow X \rightarrow B$, together with a pair of simple branch points $b', b'' \in B$ near b , each having as monodromy the transposition exchanging the sheets of $\tilde{C}$ near p and q .

In effect, we have simply introduced two new simple branch points to the cover $C \rightarrow B$, with assigned (though necessarily equal) monodromy. Note

that we can apply this construction repeatedly, to introduce any number of (pairs of) additional branch points with assigned (simple) monodromy; or we could carry out a more general construction with a number of curves C_0 .

3 Proof of the main theorem

3.1 The proof in case $B = \mathbb{P}^1$

We are now more than amply equipped to prove the theorem. We start with a morphism $\pi : X \rightarrow B$ as in 2.1. To begin with, by hypothesis X is projective; embed in a projective space and take the intersection with $\dim(X) - 1$ general hyperplanes to arrive at a smooth curve $C \subset X$. This is the curve we will start with.

What do C and the associated map $\mu : C \hookrightarrow X \rightarrow B$ look like? To answer this, start with the simplest case: suppose that the fibers X_b of π do not have multiple components, or in other words that the singular locus π_{sing} of the map π has codimension 2 in X . In this case we are done: C will miss π_{sing} altogether, so that all ramification of $\mu : C \rightarrow B$ will occur at smooth points of fibers; and simple dimension counts show that the branching will be simple. In other words, C will be pre-flexible already.

The problems start if π has multiple components of fibers. If $Z \subset X_b$ is such a component, then each point $p \in C \cap Z$ will be a ramification point of μ , and no deformation of C will move the corresponding branch point $\pi(p) \in B$. The curve C can not be flexible. And of course it's worse if π has a multiple (that is, everywhere-nonreduced) fiber: in that case π cannot possibly have a section.

To keep track of such points, let $M \subset B$ be the locus of points such that the fiber X_b has a multiple component. Outside of M , the map $\mu : C \rightarrow B$ is simply branched, and all ramification occurs at smooth points of fibers of π .

Now here is what we're going to do. First, pick a base point $p_0 \in B$, and draw a cut system: that is, a collection of real arcs joining p_0 to the branch points $M \cup N$ of μ , disjoint except at p_0 . The inverse image in C of the complement U of these arcs is simply d disjoint copies of U ; call the set of sheets Γ (or, if you prefer, label them with the integers 1 through d). Now, for each point $b \in M$, denote the monodromy around the point b by σ_b , and

express this permutation of Γ as a product of transpositions:

$$\sigma_b = \tau_{b,1}\tau_{b,2} \dots \tau_{b,k_b}$$

so that in other words

$$\tau_{b,k_b} \dots \tau_{b,2}\tau_{b,1}\sigma_b = I$$

is the identity. For future reference, let $k = \sum k_b$. We will proceed in three stages.


Stage 1: We use our second construction to produce a new curve $\tilde{C}$ with, in a neighborhood of each point $b \in M$, k_b new pairs of simple branch points $s_{b,i}, t_{b,i} \in B$, with the monodromy around $s_{b,i}$ and $t_{b,i}$ equal to $\tau_{b,i}$. Note that $\tilde{C}$ will have genus $g(C) + k$, and that the branch divisor of the projection $\tilde{\mu} : \tilde{C} \rightarrow B$ will be the union of a deformation $\tilde{N}$ of N , the points $s_{b,i}$ and $t_{b,i}$, and M . In particular we can find disjoint discs $\Delta_b \subset B$, with Δ_b containing the points b and $t_{b,1}, t_{b,2}, \dots, t_{b,k_b}$, so that the monodromy around the boundary $\partial\Delta_b$ of Δ_b is trivial.

Now, for any fixed integer n this construction can be carried out so that the curve $\tilde{C}$ has the property that $H^1(\tilde{C}, N_{\tilde{C}/X}(-q_1 - \dots - q_n)) = 0$ for any n points $q_i \in \tilde{C}$. Here we want to choose

$$n = \#N + 2k$$

so that there are global sections of the normal bundle $N_{\tilde{C}/X}$ with arbitrarily assigned values on the ramification points of $\tilde{C}$ over N and the points $s_{b,i}$ and $t_{b,i}$. This means in particular that we can deform the curve $\tilde{C}$ so as to deform the branch points of $\tilde{\mu}$ outside of M independently. What we will do, then, is

Stage 2: We will vary $\tilde{C}$ so as to *keep all the branch points $b \in N$ and all the points $s_{b,i}$ fixed*; and for each $b \in M$ specialize all the branch points $t_{b,i}$ to b within the disc Δ_b .



To say this more precisely, let $\beta \in N_1(X)$ be the class of the curve $\tilde{C}$, and consider the maps

$$\overline{\mathcal{M}}_{g',0}(X, \beta) \longrightarrow \overline{\mathcal{M}}_{g',0}(B, d) \longrightarrow B_{2d+2g'-2}$$

with the second map assigning to a stable map $C \rightarrow B$ its branch divisor. What we are saying is, starting at the branch divisor

$$D_1 = \tilde{N} + \sum s_{b,i} + \sum t_{b,i} + \sum_{b \in M} k_b \cdot b$$

of the map $\tilde{\mu}$, draw an analytic arc $\gamma = \{D_\lambda\}$ in the subvariety

$$\Phi = \tilde{N} + \sum s_{b,i} + \sum_{b \in M} k_b \cdot b + \sum (\Delta_b)_{k_b} \subset B_{2d+2g'-2}$$

tending to the point

$$D_0 = \tilde{N} + \sum s_{b,i} + 2 \sum_{b \in M} k_b \cdot b.$$

Since the image of the composition

$$\overline{M}_{g',0}(X, \beta) \longrightarrow B_{2d+2g'-2}$$

contains Φ , we can find an arc $\delta = \{f_\nu\}$ in $\overline{M}_{g',0}(X, \beta)$ that maps onto γ , with f_1 the inclusion $\tilde{C} \hookrightarrow X$.

Stage 3: Let $f_0 : C_0 \rightarrow X$ be the limit, in $\overline{M}_{g',0}(X, \beta)$, of the family of curves constructed in Stage 2: that is, the point of the arc δ over $D_0 \in \Phi \subset B_{2d+2g'-2}$. Let $A \subset C_0$ be the normalization of any irreducible component of C_0 on which the composition πf_0 is nonconstant (that is, whose image is not contained in a fiber), and let $f : A \rightarrow X$ be the restriction of f_0 to A .

By construction, the composition πf is unramified over a neighborhood of M : the monodromy around the boundary $\partial\Delta_b$ of each disc Δ_b is trivial, and it can be branched over at most one point b inside Δ_b , so it can't be branched at all over Δ_b . Indeed, it is (at most) simply branched over each point of N and each point $s_{b,i}$, and unramified elsewhere. Moreover, since we can carry out the specialization of $\tilde{C}$ above with the entire fiber of $\tilde{C}$ over the points of N and the $s_{b,i}$ fixed, the ramification of πf on A over these points will occur at smooth points of the corresponding fibers of π . In other words, *the map $f : A \rightarrow X$ is preflexible*, and we are done. □

3.2 The proof for arbitrary curves B

As we indicated at the outset, there are two straightforward ways of extending this result to the case of arbitrary curves B .

For one thing, virtually all of the argument we have made goes over without change to the case of base curves B of any genus h . The one exception to this is the statement that the space $\overline{M}_{g,0}(B, d)$ of stable maps $f : C \rightarrow B$ of degree d from curves C of genus g to B has a unique irreducible component whose general member corresponds to a flat map $f : C \rightarrow B$. This is false in general—consider for example the case $g = d(h - 1) + 1$ of unramified covers. It is true, however, if we restrict ourselves to the case $g \gg h, d$ (that is, we have a large number of branch points) and look only at covers whose monodromy is the full symmetric group S_d . Given this fact, and observing that our second construction allows us to increase the number of branch points of our covers $C \rightarrow B$ arbitrarily, the theorem can be proved for general B just as it is proved above for $B \cong \mathbb{P}^1$.

Alternatively, Johan deJong showed us a simple way to deduce the theorem for general B from the case $B \cong \mathbb{P}^1$ alone. We argue as follows: given a map $\pi : X \rightarrow B$ with rationally connected general fiber, we choose any map $g : B \rightarrow \mathbb{P}^1$ expressing B as a branched cover of $\mathbb{P}^1$. We can then form the “norm” of X : this is the (birational isomorphism class of) variety $Y \rightarrow \mathbb{P}^1$ whose fiber over a general point $p \in \mathbb{P}^1$ is the product

$$Y_p = \prod_{q \in g^{-1}(p)} X_q.$$

Since the product of rationally connected varieties is again rationally connected, it follows from the $\mathbb{P}^1$ case of the theorem that $Y \rightarrow \mathbb{P}^1$ has a rational section, and hence so does π .

4 An example

There are a number of disquieting aspects of the argument in Section 3.1, and in particular about the specialization in Stage 2 of that argument. Clearly the curve $f : A \rightarrow X$ constructed there cannot meet any multiple component of a fiber of $\pi : X \rightarrow B$; that is, for each $b \in M$ it must meet the fiber X_b only in reduced components of X_b . This raises a number of questions: what if the fiber X_b is multiple? How can the curve $\tilde{C}$, which meets all the multiple components of X_b , specialize to one that misses them all? And can we say which reduced components of X_b the curve A will meet?

The answers to the first two questions are straightforward: in fact, the argument given here proved that *the map $\pi : X \rightarrow B$ cannot have multiple fibers*, that is, every fiber X_b must have a reduced component.¹ As for the second, what must happen is that as our parameter $\delta \rightarrow 0$, the points of intersection of C_δ with the multiple components of X_b slide toward the reduced components of X_b ; the curve C_0 produced in the limit will have components contained in the fiber X_b and joining the points of intersection of A with X_b to each of the multiple components. Finally, the answer to the third question—and indeed the whole process—may be illuminated by looking at a simple example; we will do this now.


To start, we have to find an example of a map $\pi : X \rightarrow B$ with rationally connected general fiber and a special fiber having a multiple component (and

¹In fact, this assertion is nearly tantamount to Theorem 1.1 itself, but we were unable to prove it directly except under additional and restrictive hypotheses

smooth total space X). Without question, the simplest example will have general fiber $X_b \cong \mathbb{P}^1$, and special fiber a chain of three smooth rational curves:



The middle component will have multiplicity 2 in the fiber, and self-intersection -1 ; the outer two components will each appear with multiplicity 1 in the fiber, and will have self-intersection -2 . The simplest way to construct a family with such a fiber is to start with a trivial family $X_0 = \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$, blow up any point p , and then blow up the point q of intersection of the exceptional divisor with the proper transform of the fiber through p to obtain X .




We will denote by F the proper transform in X of the fiber through p in

$\mathbb{P}^1 \times \mathbb{P}^1$, and by G the proper transform of the first exceptional divisor; the second exceptional divisor—the multiple component of the special fiber—we will call E . To arrive at the simplest possible curve $\tilde{C} \subset X$ meeting the multiple component E of the special fiber of this family, we start with a curve $C \subset \mathbb{P}^1 \times \mathbb{P}^1$ of degree 2 over B that is simply tangent to the special fiber at the point p ; the proper transform $\tilde{C}$ of C in X will then meet E once transversely and F and G not at all. (We’re not trying to make excuses here, but note that it’s virtually impossible to draw a decent picture of the configuration $\tilde{C} \subset X \rightarrow B$: the curve $\tilde{C}$ is supposed to meet E once transversely, but still have degree 2 over B and be ramified over B at its point of intersection with E .)

Now that we’ve got this set up, what happens when we push another branch point of $\tilde{C} \rightarrow B$ in to the special fiber of π ? The answer is that one of three things can happen, two generically. We will describe these first geometrically in terms of the original curve $C \subset \mathbb{P}^1 \times \mathbb{P}^1$ and its proper transforms, and then write down typical equations.

One possibility is that the ramification point p of C over B becomes a node. In this case the limit C_0 of the proper transforms C_ν of the curves will actually contain the component E of the special fiber (the limit of the proper transforms is not the proper transform of the limiting curve, but rather its total transform minus the divisor $E + G$). The remaining component—the actual proper transform of the limiting curve—will have two distinct sheets in a neighborhood of the special fiber, meeting G transversely in distinct points, and each of course unramified over B :




This specialization is easy to see in terms of equations: if we choose affine coordinates x on our base $\mathbb{P}^1$ and y on the fiber, we can write the equation of our family $\{C_\nu\}$ of curves as


$$C_\nu : y^2 = x^2 - \nu x$$

and specialize the branch point over $x = \nu$ simply by letting $\nu \rightarrow 0$. We can see either from this family of equations, or geometrically, that as ν tends to 0 the point of intersection of the proper transform $\tilde{C}_\nu$ of C_ν slides along E toward the point of intersection $E \cap G$; when it reaches $E \cap G$ the limiting curve becomes reducible, splitting off a copy of G .

Now, by the symmetry of $X \rightarrow B$ —we could also blow down the curves E and F in X to obtain $\mathbb{P}^1 \times \mathbb{P}^1$ —we would expect that there would be a similar specialization with the roles of F and G reversed, and there is: if the curve $C \subset \mathbb{P}^1 \times \mathbb{P}^1$ specializes to one containing the fiber $\{0\} \times \mathbb{P}^1$, the limit $\tilde{C}_0$ of the proper transforms will (generically) consist of the union of F with a curve A , with A unramified of degree 2 over B in a neighborhood of the special fiber and meeting the special fiber in two distinct points of F .



Finally, there is a common specialization of these two families: if the curve $C \subset \mathbb{P}^1 \times \mathbb{P}^1$ specializes to one that both contains the fiber $\{0\} \times \mathbb{P}^1$ and is singular at the point q —that is, consists in a neighborhood of the special fiber of the fiber and two sections, one passing through p —then the limit $\tilde{C}_0$ of the proper transforms will consist of the union of all three components E , F and G of the special fiber with a curve A consisting of a section meeting the special fiber in a point of F and a section meeting the special fiber in a point of E :



It's also very instructive to look at this example from the point of view of the equations of the curves. To begin with, denote by $|\mathcal{O}_X(d, e)|$ the total transform of the linear system of curves of bidegree (d, e) on $\mathbb{P}^1 \times \mathbb{P}^1$. We are looking here at the linear system

$$\tilde{\mathcal{D}} = |\mathcal{O}_X(1, 2)(-G - 2E)|,$$

that is, the proper transform of the linear series $\mathcal{D}$ of curves $C \subset \mathbb{P}^1 \times \mathbb{P}^1$ of bidegree $(1, 2)$ that pass through p with vertical tangent. Explicitly, these curves form a 3-dimensional linear series, which we may write in affine coordinates (x, y) on $\mathbb{P}^1 \times \mathbb{P}^1$ as

$$\mathcal{D} = \{axy^2 + by^2 + cxy + dx\}_{[a,b,c,d] \in \mathbb{P}^3}$$

Writing the equation of a typical member of $\mathcal{D}$ as a polynomial in y :

$$(ax + b) \cdot y^2 + (cx) \cdot y + (dx) = 0$$

we see that its branch divisor is the zero locus of the quadratic polynomial

$$(dx)^2 - 4(ax + b)(dx) = (c^2 - 4ad) \cdot x^2 - 4db \cdot x$$

whose roots are at $x = 0$ and $x = 4db/(c^2 - 4ad)$. It's probably best to express this in terms of the maps

$$\overline{M}_{0,0}(X, [\tilde{\mathcal{D}}]) \longrightarrow \overline{M}_{0,0}(B, 2) \longrightarrow B_2$$

introduced in section 3 above. Here, the variety $\overline{M}_{0,0}(X, [\tilde{\mathcal{D}}])$ has a component M which is a blow-up of the $\mathbb{P}^3_{[a,b,c,d]}$ parametrizing the linear series $\tilde{\mathcal{D}}$ (it also has a second, extraneous component whose general point corresponds to a map $f : C \rightarrow X$ with reducible domain and image containing the line $y = 0$ doubly; this component is not involved here). The image of the composite map $M \rightarrow B_2$ is simply the locus $B_0 \cong B \subset B_2$ of divisors of degree 2 in $B \cong \mathbb{P}^1$ containing the point $x = 0$, with the map

$$\eta : M \rightarrow \mathbb{P}^3_{[a,b,c,d]} \rightarrow B_0 \cong \mathbb{P}^1$$

given by

$$[a, b, c, d] \mapsto [4db, c^2 - 4ad].$$

What we see in particular from this is that *the fiber of η over the point $x = 0$ is reducible*, with components given by $d = 0$ and $b = 0$. Now, in Stage 2 of our argument, as applied here, we start with an arc $\gamma \subset B_0 \subset B_2$ in which the second branch point approaches $x = 0$, and lift that to an arc $\delta \subset M$. If our arc $\delta \subset M$ lifting the arc $\gamma \subset B_0 \subset B_2$ approaches the component $d = 0$ —whose general member corresponds to a curve $C \subset \mathbb{P}^1 \times \mathbb{P}^1$ singular at p —we get a family of stable maps whose limit is as described in the first example above. If, on the other hand, it approaches the component $b = 0$, whose general member corresponds to a curve $C \subset \mathbb{P}^1 \times \mathbb{P}^1$ containing the fiber $x = 0$, we get a limit as depicted in the third example. And finally, if δ approaches (generically) a point in the intersection of these two components, we get an example of the third type.

References

- [Ca] F. Campana, *Connexité rationnelle des variétés de Fano*, Ann. Sc. E. N.S. **25** (1992), 539-545
- [C] A. Clebsch, *Zur Theorie der Riemann'schen Flächen*, Math Ann. **6** (1872), 216-230 Springer-Verlag, Berlin, 1996.

- [H] A. Hurwitz, *Ueber Riemann'sche Flächen mit gegebenen Verzweigungspunkten*, Math. Ann. **39** (1891) 1-61.
- [K] J. Kollár, *Rational Curves on Algebraic Varieties*, Ergebnisse der Math. 32, Springer-Verlag, Berlin, 1996.
- [KMM] J. Kollár, Y. Miyaoka, S. Mori, *Rationally Connected Varieties*, J. Alg. Geom. **1** (1992) 429-448